

Super 8 Mineral with ClariFly®

Product number: 66082

A nutritional supplement containing Diflubenzuron Insect Growth Regulator for continuous feeding to cattle through the fly season. Prevents the emergence of horn flies, face flies, house flies, and stable flies from manure of treated cattle.

Product Info

When used as directed, Stockade® Super 8 Mineral ClariFly® provides sufficient diflubenzuron insect growth regulator to prevent the emergence of adult house flies, stable flies, face flies, and horn flies from manure of treated cattle. Existing adult flies will not be affected. Start feeding before flies appear and continue use until cold weather marks the end of the fly season.



- Excellent all-around mineral for cattle, horses or goats on good to excellent quality forages and/or those receiving additional fortification from other feed sources.
- Contains 8% Phosphorus, and other macro and trace minerals along with vitamins essential for the animal's overall diet
- This mineral can be easily mixed and matched with any of the STOCKADE protein supplement blocks or tubs to create a custom supplement program.
- Added ClariFly prevents the emergence of horse, face, house and stable flies from manure of treated cattle. For best results, feed continuously through fly season.

Nutrient Guarantee

Active Ingredient Concentration	Diflubenzuron* (CAS# 35367-38-5) 0.04%
Calcium, Min	19.00%
Calcium, Max	22.80%
Phosphorus, Min	8.00%
Salt, Min	10.00%
Salt, Max	12.00%
Copper, Min	250 ppm
Copper, Max	270 ppm
Iodine, Min	100 ppm
Manganese, Min	1,000 ppm
Selenium, Min	26 ppm
Zinc, Min	800 ppm
Vitamin A, Min	50,000 IU/lb
Vitamin D-3, Min	5,000 IU/lb
Vitamin E, Min	50 IU/lb

Product Ingredients

Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Molasses Products, Processed Grain By-Products, Vegetable Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Selenium Yeast, Brewer's Dried Yeast, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil, Artificial Flavoring and Red Iron Oxide.

Product directions

Place Stockade® Super 8 Mineral ClariFly® in feeders close to water sources and/or resting areas. Allow free choice to cattle. Cattle should consume an average of 0.4 ounces of Stockade® Super 8 Mineral ClariFly® per 100 lbs. of animal body weight per day, which is equivalent to 4 ounces per day for a 1,000 pound animal.

1. Assure cattle are not starved for nutrients by providing them prior to feeding this product.
2. This product contains added copper. Do not feed to sheep or other species that have a low tolerance to supplemental copper.
3. Offer cattle 50 pounds of this product for every 15 head of cattle.
4. Remove all other self-limiting nutritional supplements.
5. Locate this product where cattle congregate (watering, loafing, shade areas).
6. Replenish this product on a regular basis as needed.
7. If intake of this product is below 0.4 oz. per 100 lbs. animal body weight per day, increase the number of feeding locations and/or change locations to areas more frequented by cattle.
8. If intake of this product is above 0.4 oz. per 100 lbs. animal body weight per day, reduce the number of feeding locations and/or change locations to areas less frequented by cattle.
9. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired.

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash clothing before reuse.

Personal Protective Equipment (PPE): Applicators and other handlers of this product must wear:

- o Long sleeve shirt and long pants
- o Chemical resistant gloves
- o Shoes
- o Socks

First Aid	
IF IN EYES	<ul style="list-style-type: none">o Hold eye open and rinse slowly and gently with water for minutes.o Remove contact lenses, if present, after the first 5 minutes; continue rinsing.o Call a poison control center or doctor for treatment advice.
IF ON SKIN	<ul style="list-style-type: none">o Take off contaminated clothing.o Rinse skin with water for 15-20 minutes.o Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control doctor or going for treatment. For additional information on this pesticide including health concerns, medical emergencies or pesticide incidents, 222-1222.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by stc disposal. **Pesticide Storage:** Keep this product in its tightly closed original

Store in a cool, dry (preferably locked) area that is inaccessible to children and animals. **Pesticide Disposal:** Wastes resulting from the use of this pesticide may be disposed of on-site or at an approved waste disposal facility. **Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into mixing equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by other procedures approved by state and local authorities.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.